

Cardiovascular Imaging In-a-Month

Transient ST Elevation During Radiofrequency Energy Application From the Left Sinus of Valsalva

Tomoya HIRATSUJI, MD
Hiroshi TADA, MD
Shigeto NAITO, MD
Shigeru OSHIMA, MD, FJCC
Koichi TANIGUCHI, MD,

FJCC




Fig. 1

群馬県立心臓血管センター 循環器内科(平辻知也, 多田 浩, 内藤滋人, 大島 茂, 谷口興一): 〒371-0004 群馬県前橋市 亀泉町甲3-12

Division of Cardiology, Gunma Prefectural Cardiovascular Center, Gunma

Address for correspondence: TADA H, MD, Division of Cardiology, Gunma Prefectural Cardiovascular Center, Ko 3-12, Kameizumi-cho, Maebashi, Gunma 371-0004

Manuscript received February 24, 2003; revised March 28, 2003; accepted April 10, 2003




Fig. 2

CASE

A 46-year-old man was referred to our center for radiofrequency catheter ablation for the treatment of symptomatic, monomorphic premature ventricular contractions (Fig. 1 - A). During an electrophysiologic study, the earliest ventricular activation was identified within the left sinus of Valsalva. After identifying the ostium of the left coronary artery and the contour of the left sinus of Valsalva by angiography (Fig. 2 - A), an ablation catheter was positioned within the left sinus of Valsalva (Fig. 2 - B). The first application of radiofrequency energy within the left sinus of Valsalva had no effect. No ST-T segment change was observed during the application. After detailed mapping toward the right sinus of Valsalva, a second application was performed within the left sinus of Valsalva where the ventricular activation recorded from the distal electrodes of the ablation catheter preceded the onset of the QRS complex by 23 msec. Ten seconds after application of radiofrequency energy using a maximum power of 35 watts and a maximum electrode-tissue interface temperature of 55 °C, ST segment elevation in the inferior leads appeared (Fig. 1 - B - left, arrows). After prompt cessation of the radiofrequency energy, the ST segment elevation in these leads returned to the baseline (Fig. 1 - B - right), and the patient did not complain of chest pain.

Points for Diagnosis

Recently, ventricular tachycardias or premature ventricular contractions originating from the left ventricular epicardial site have been ablated from the aortic sinus of Valsalva^{1,2}). The 12-lead electrocardiogram of this kind of arrhythmia often shows an S wave in lead V_1 , tall R waves in the inferior leads, precordial R wave transition in leads $V_1 - V_3$, and no S wave in either lead V_5 or V_6 like this case (**Fig. 1**).

In this case, selective right coronary angiography revealed an anomalous origin of the right coronary artery from the left sinus of Valsalva, close to the right sinus of Valsalva (**Fig. 3**, arrows). The ablation site where the earliest ventricular activation was recorded was at the ostium of the right coronary artery. Radiofrequency energy application at the ostium might cause spasm of the right coronary artery. The patient was discharged under medication with 300 mg/day of mexiletine and has not complained of chest pain during a 9-month follow-up period.

Coronary anomalies affect about 1% of the general population³). Anomalous origin of the right coronary artery from the left sinus of Valsalva is one of the most common coronary anomalies⁴). Selective coronary angiography of both coronary arteries should be performed before radiofrequency

application to avoid potential complications in patients undergoing radiofrequency application from the aortic sinus.

Diagnosis: Anomalous origin of the right coronary artery from the left sinus of Valsalva

Key Words: Heart defects, congenital (anomalous origin of the coronary artery); Ablation-catheter ; Ventricular arrhythmia

References

- 1) Kanagaratnam L, Tomassoni G, Schweikert R, Pavia S, Bash D, Beheiry S, Niebauer M, Saliba W, Chung M, Tchou P, Natale A: Ventricular tachycardias arising from the aortic sinus of Valsalva: An under-recognized variant of left outflow tract ventricular tachycardia. *J Am Coll Cardiol* 2001; **37**: 1408 - 1414
- 2) Ouyang F, Fotuhi P, Ho SY, Hebe J, Volkmer M, Goya M, Burns M, Antz M, Ernst S, Cappato R, Kuck KH: Repetitive monomorphic ventricular tachycardia originating from the aortic sinus cusp: Electrocardiographic characterization for guiding catheter ablation. *J Am Coll Cardiol* 2002; **39**: 500 - 508
- 3) Angelini P, Velasco JA, Flamm S: Coronary anomalies: Incidence, pathophysiology, and clinical relevance. *Circulation* 2002; **105**: 2449 - 2454
- 4) Ho JS, Strickman NE: Anomalous origin of the right coronary artery from the left coronary sinus: Case report and literature review. *Tex Heart Inst J* 2002; **29**: 37 - 39




Fig. 3

Fig. 1 Twelve-lead electrocardiograms

A: Twelve-lead electrocardiogram showing premature ventricular contraction as an S wave in lead I, tall R waves in the inferior leads, precordial R wave transition in leads V₂-V₃, and no S wave in either lead V₅ or V₆.

B: Twelve-lead electrocardiogram during and just after radiofrequency energy application from the left sinus of Valsalva showing ST segment elevation in the inferior leads during radiofrequency energy application (left, arrows), which returned to the baseline just after cessation of the radiofrequency energy (right).

RF = radiofrequency.

Fig. 2 Angiograms obtained in the right anterior oblique (RAO 35°, upper row) and left anterior oblique

(LAO 45°, lower row) projections

A: Left coronary angiogram showing the ostium of the left coronary artery and the shape of the left sinus of Valsalva.

B: Angiogram of the ablation site showing a 7 F quadripolar catheter with a 4 mm distal electrode (Biosense Webster, Diamond Bar) positioned within the left sinus of Valsalva (arrows).

Fig. 3 Angiograms obtained in the right anterior oblique (RAO 35°, left) and left anterior oblique (LAO 45°, right) projections

Right coronary angiogram showing the origin from the left sinus of Valsalva, close to the right sinus of Valsalva. The ablation site where the earliest ventricular activation was recorded was at the ostium of the right coronary artery (arrows).